

DUALITY OF MULTITIME VECTOR INTEGRAL PROGRAMMING WITH QUASIINVEXITY

ȘTEFAN MITITELU, VASILE PREDA AND MIHAI POSTOLACHE

ABSTRACT. We consider a multitime vector variational problem (VVP) and a multitime vector fractional variational problem (VFP). In their recently published work [Balkan J. Geom. Appl., **16**(2011), No. 2, 90-101], Mititelu and Postolache established necessary efficiency conditions for the two problems. By means of these conditions and using the framework of (ρ, b) -quasiinvexity, in this work we introduce duality conditions of Mond-Weir-Zalmai type for the fractional problem (VFP) through weak, direct and converse duality theorems. Our approach allows us to obtain some duality conditions for (VVP) as special cases in this theory.

1. INTRODUCTION

Beginning with the research of Valentine [23], published in 1937, during the years, the variational problem with constraints known different steps in its development. In time, several authors have been interested in the study of (sufficient) efficiency conditions and duality for fractional vector programs in connection with generalized convexity. To quote some sources, see [6] by Jeyakumar and Mond, [7] by Khan and Hanson, [8] by Liang et al., [10] by Mititelu, [15] and [16] by Pitea et al., [19] by Reddy and Mukherjee, [20] by Singh and Hanson.

In 2007 Mititelu and Stancu-Minasian [12] considered the following vector (or multiobjective) fractional variational problem:

$$(MSP) \quad \left\{ \begin{array}{l} \text{Maximize} \quad \left(\frac{\int_a^b f_1(t, x, \dot{x}) dt}{\int_a^b k_1(t, x, \dot{x}) dt}, \dots, \frac{\int_a^b f_p(t, x, \dot{x}) dt}{\int_a^b k_p(t, x, \dot{x}) dt} \right) \\ \text{subject to} \quad x(a) = a_0, \quad x(b) = b_0, \\ g(t, x, \dot{x}) \leq 0, \quad h(t, x, \dot{x}) = 0, \quad \forall t \in I, \end{array} \right.$$

Received: December 20, 2010. *Revised:* June 10, 2011.

2010 *Mathematics Subject Classification:* 65K10, 90C29, 26B25.

Key words and phrases: Multitime vector fractional variational problem, efficient solution, normal efficient solution, (ρ, b) -quasiinvexity.

where $I = [a, b]$ is interval, $x = (x_1, \dots, x_n): I \rightarrow \mathbb{R}^n$ is piecewise smooth function on I with $\dot{x} = \frac{dx}{dt}$ its derivative, $f_1, k_1, \dots, f_p, k_p: I \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$, $g: I \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $h: I \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^q$ are functions of C^2 -class.

Mititelu [9], Mititelu and Stancu-Minasian [12] established necessary efficiency (Pareto minimum) conditions for problem (MSP). Using generalized quasiinvex functions, they also developed a dual program theory stating weak, direct and converse duality theorems [12].

In 2006, Udriște [21] studied a control variational problem within multitime framework, establishing some optimality conditions, that is he introduced a multitime maximum principle for this kind of problems. Meanwhile, Udriște and Țevy [22] established the multitime principle of maximum for a multitime variational problem. In 2009 Pitea, Udriște and Mititelu [15], [16] considered the multitime vector variant of the problem (MSP) in geometrical language. Using curvilinear integrals they established necessary efficiency conditions and developed a duality theory for this problem. Recently, Mititelu and Postolache [11] studied the same objectives for multitime vector fractional and nonfractional variational problems on Riemannian manifolds, but using multiple integrals on a measurable set Ω . For a survey of recent developments in multiobjective optimization, we address the reader to [1], by Chinchuluun and Pardalos.

It is our purpose in the present paper to introduce necessary efficiency conditions for multitime vector fractional and nonfractional variational problems involving also multiple integrals on a measurable set Ω , but in the space \mathbb{R}^n . For these problems we develop some duality results through weak, direct and converse theorems.

Consider a measurable set Ω in \mathbb{R}^m and the functions

$$\begin{aligned} x: \Omega &\rightarrow \mathbb{R}^n, & X: \Omega \times \mathbb{R}^n \times \mathbb{R}^{nm} &\rightarrow \mathbb{R}, \\ f = (f_r): \Omega \times \mathbb{R}^n \times \mathbb{R}^{nm} &\rightarrow \mathbb{R}^p, & k = (k_r): \Omega \times \mathbb{R}^n \times \mathbb{R}^{nm} &\rightarrow \mathbb{R}^p, \\ g = (g_\alpha): \Omega \times \mathbb{R}^n \times \mathbb{R}^{nm} &\rightarrow \mathbb{R}^m, & h = (h_\beta): \Omega \times \mathbb{R}^n \times \mathbb{R}^{nm} &\rightarrow \mathbb{R}^q, \end{aligned}$$

where $m, q \in \mathbb{N}^*$, $r = \overline{1, n}$, $\alpha = \overline{1, m}$ and $\beta = \overline{1, q}$. Denote by $t = (t^1, \dots, t^m) = (t^\alpha)$ the current element of Ω and by $x = (x^1, \dots, x^n) = (x^k)$ the current element of \mathbb{R}^n . The arguments of $X, f, k, g_\alpha, h_\beta$ are $(t, x, x_\gamma) = (t, x(t), x_\gamma(t))$, where

$$x = x(t) = (x^1(t), \dots, x^n(t)) = (x^k(t)), \quad t \in \Omega$$

and

$$x_\gamma = x_\gamma(t) = \frac{\partial x}{\partial t^\gamma}(t), \quad \gamma = \overline{1, m}$$

is the first derivative of $x = x(t)$. We suppose that $X, f, k, g_\alpha, h_\beta \in C^2(\Omega)$.

In this paper, Euler-Lagrange equations of the form

$$\frac{\partial X}{\partial x^k} - \frac{\partial}{\partial t^1} \left(\frac{\partial X}{\partial x_{t^1}^k} \right) - \dots - \frac{\partial}{\partial t^m} \left(\frac{\partial X}{\partial x_{t^m}^k} \right) = 0, \quad k = \overline{1, n}$$

are shortly written as

$$\frac{\partial X}{\partial x^k} - \frac{\partial}{\partial t^\gamma} \left(\frac{\partial X}{\partial x_\gamma^k} \right) = 0, \tag{1.1}$$

where we denoted $x_\gamma^k = \frac{\partial x^k}{\partial t^\gamma}$ and

$$\begin{aligned} \frac{\partial}{\partial t^\gamma} \left(\frac{\partial X}{\partial x_\gamma^k} \right) &= \frac{\partial^2 X}{\partial t^\gamma \partial x_\gamma^k} + \frac{\partial^2 X}{\partial x^i \partial x_\gamma^k} \frac{\partial x^i}{\partial t^\gamma} + \frac{\partial^2 X}{\partial x_s^i \partial x_\gamma^k} \frac{\partial x_s^i}{\partial t^\gamma} \\ &= \frac{\partial^2 X}{\partial t^\gamma \partial x_\gamma^k} + \frac{\partial^2 X}{\partial x^i \partial x_\gamma^k} \frac{\partial x^i}{\partial t^\gamma} + \frac{\partial^2 X}{\partial x_s^i \partial x_\gamma^k} \frac{\partial^2 x^i}{\partial t^s \partial t^\gamma}. \end{aligned}$$

In the following, we use the Euler-Ostrogradsky equations with the vector form given by (1.1), namely

$$\frac{\partial X}{\partial x} - \frac{\partial}{\partial t^\gamma} \left(\frac{\partial X}{\partial x_\gamma} \right) = 0.$$

We define the set $\Phi = \{x: \Omega \rightarrow M \mid x \text{ is piecewise smooth on } \Omega\}$, where Ω is a normed space with $\|x\| = \|x\|_\infty + \sum_{k=1}^n \|\dot{x}^k\|_\infty$.

Throughout the paper, for two vectors $v = (v_1, \dots, v_n)$ and $w = (w_1, \dots, w_n)$ the relations of the form $v = w$, $v < w$, $v \leq w$, $v \leq w$ are defined as follows

$$\begin{aligned} v = w &\Leftrightarrow v_i = w_i, \quad i = \overline{1, n}; & v < w &\Leftrightarrow v_i < w_i, \quad i = \overline{1, n}; \\ v \leq w &\Leftrightarrow v_i \leq w_i, \quad i = \overline{1, n}; & v \leq w &\Leftrightarrow v \leq w \text{ and } v \neq w. \end{aligned}$$

Important note. To simplify the presentation of this work, in our subsequent theory, we shall set

$$\pi_x(t) = (t, x(t), x_\gamma(t)), \quad \pi_{x^0}(t) = (t, x^0(t), x_\gamma^0(t)), \quad \pi_y(t) = (t, y(t), y_\gamma(t)).$$

On X , consider the functionals

$$F_r(x) = \int_\Omega f_r(\pi_x(t))dv, \quad K_r(x) = \int_\Omega k_r(\pi_x(t))dv, \quad r = \overline{1, p},$$

and assume $K_r(x) \neq 0$, $r = \overline{1, p}$.

In this paper, we establish various conditions of Mond-Weir duality type for the following multitime vector fractional variational problem:

$$(VFP) \begin{cases} \text{Maximize} & \left(\frac{F_1(x)}{K_1(x)}, \dots, \frac{F_p(x)}{K_p(x)} \right) \\ \text{subject to} & g_\alpha(\pi_x(t)) = 0, \quad h_\beta(\pi_x(t)) \leq 0, \\ & x(t)|_{\partial\Omega} = u(t) \text{ (given)}, \forall t \in \Omega, \end{cases}$$

where the fractions are well defined and $dv = dt_1 dt_2 \dots dt_n$.

In particular, there are obtained duality conditions for the following multitime vector nonfractional variational problem

$$(VVP) \quad \begin{cases} \text{Maximize } I[x] = \int_{\Omega} f(\pi_x(t))dv \\ \text{subject to } g_{\alpha}(\pi_x(t)) = 0, \quad h_{\beta}(\pi_x(t)) \leq 0, \\ x|_{\partial\Omega} = u(t), \forall t \in \Omega, \end{cases}$$

The two problems, stated above, have the same domain

$$\mathcal{D} = \{x \in \Phi \mid g_{\alpha}(\pi_x(t)) = 0, \quad h_{\beta}(\pi_x(t)) \leq 0, \quad x|_{\partial\Omega} = u(t), \quad \forall t \in \Omega\}.$$

2. NECESSARY EFFICIENCY CONDITIONS FOR PROBLEMS (VVP) AND (VFP)

In their joint paper [11], Mititelu and Postolache established necessary efficiency conditions in the geometric framework of a Riemannian manifold for problems (VVP) and (VFP). In what follows, we shall present these conditions adapted to the study in the real space \mathbb{R}^n .

A. Efficiency for multitime vector variational problem (VVP). The vector functional

$$I[x] = \int_{\Omega} f(\pi_x(t))dv,$$

can be written componentwise as

$$I[x] = (F_1(x), \dots, F_p(x)).$$

Definition 2.1 ([3]). A point $x^* \in \mathcal{D}$ is said to be *efficient solution* (Pareto minimum) to (VVP) if there exist no $x \in \mathcal{D}$ such that $I[x] \leq I[x^*]$.

Now, according to [11], in the \mathbb{R}^n case we have the following necessary efficiency conditions for (VVP).

Theorem 2.1 (Necessary efficiency for (VVP)). *Consider the vector multitime variational problem (VVP) and let $x = x(t) \in \mathcal{D}$ be an efficiency solution to (VVP). Then there are vector functions $\tau \in \mathbb{R}^p$, $\lambda(t) \in \mathbb{R}^m$ and $\mu(t) \in \mathbb{R}^q$, all being piecewise smooth functions, which satisfy the conditions*

$$(VFJ) \quad \begin{cases} \tau^r \frac{\partial f_r}{\partial x^k} + \lambda^{\alpha}(t) \frac{\partial g_{\alpha}}{\partial x^k} + \mu^{\beta}(t) \frac{\partial h_{\beta}}{\partial x^k} - \\ \quad - \frac{\partial}{\partial t^{\gamma}} \left(\tau^r \frac{\partial f_r}{\partial x^{\gamma k}} + \lambda^{\alpha}(t) \frac{\partial g_{\alpha}}{\partial x^{\gamma k}} + \mu^{\beta}(t) \frac{\partial h_{\beta}}{\partial x^{\gamma k}} \right) = 0, \\ \mu^{\beta}(t) h_{\beta}(\pi_x(t)) = 0 \text{ (no summation), } \beta = \overline{1, q}, \\ \tau = (\tau^r) \geq 0, \quad (\mu^{\beta}(t)) \geq 0, \quad t \in \Omega, \quad \gamma = \overline{1, m}, \quad k = \overline{1, n}. \end{cases}$$

Definition 2.2 ([9]). The efficient solution $x^0 \in \mathcal{D}$ is called *normal efficient solution* to (VVP) if conditions (VFJ) are satisfied with

$$\tau \geq 0, \quad e'\tau = 1, \quad e = (1, \dots, 1) \in \mathbb{R}^p,$$

where $'$ is the transposition sign.

B. Efficiency for multitime vector fractional variational problem (VFP).

In this section, we present the necessary efficiency conditions for (VFP) established in [11], also adapted to the study in \mathbb{R}^n . Remark that the definition of efficient solution for (VFP) is similar Definition 2.1.

Theorem 2.2 (Necessary efficiency for (VFP)). *Let $x = x(t) \in \mathcal{D}$ be a normal efficient solution to problem (VFP). Then there exist vector $\tau = (\tau^r) \in \mathbb{R}^p$, piecewise smooth functions $\lambda = (\lambda^\alpha(t)) \in \mathbb{R}^{nm}$ and $\mu = (\mu^\beta(t)) \in \mathbb{R}^q$ that satisfy the conditions*

$$(MFJ) \quad \begin{cases} \tau^r \left[K_r(x) \frac{\partial f_r}{\partial x^k} - F_r(x) \frac{\partial k_r}{\partial x^k} \right] + \lambda^\alpha(t) \frac{\partial g_\alpha}{\partial x^k} + \mu^\beta(t) \frac{\partial h_\beta}{\partial x^k} - \\ - \frac{\partial}{\partial t^\gamma} \left\{ \tau^r \left[K_r(x) \frac{\partial f_r}{\partial x_\gamma^k} - F_r(x) \frac{\partial k_r}{\partial x_\gamma^k} \right] + \lambda^\alpha(t) \frac{\partial g_\alpha}{\partial x_\gamma^k} + \mu^\beta(t) \frac{\partial h_\beta}{\partial x_\gamma^k} \right\} = 0, \\ \mu^\beta(t) h_\beta(\pi_x(t)) = 0 \text{ no summation, } \beta = \overline{1, q}, \\ \tau \geq 0, \quad e'\tau = 1, \quad (\mu^\beta(t)) \geq 0, \quad t \in \Omega, \quad \gamma = \overline{1, m}, \quad k = \overline{1, n}. \end{cases}$$

The definition of normal efficient solution for (VFP) is similar Definition 2.2.

Let ρ be a real number, and $b: \Phi \times \Phi \rightarrow [0, \infty)$ a nonnegative functional. On Φ , consider the functional

$$\mathcal{X}(x) = \int_{\Omega} X(\pi_x(t)) dv.$$

Definition 2.3 ([21]). The functional \mathcal{X} is said to be (strictly) (ρ, b) -quasiinvex at x^0 if there exist vector functions $\eta(t) = (\eta_1(t), \dots, \eta_m(t)) \in \mathbb{R}^n$ of C^1 -class with $\eta(t)|_{\partial\Omega} = 0$ and $\theta(x, x^0) \in \mathbb{R}^n$ such that for any x ($x \neq x^0$),

$$\begin{aligned} \mathcal{X}(x) &\leq \mathcal{X}(x^0) \Rightarrow \\ b(x, x^0) \int_{\Omega} \left\{ \eta_k \frac{\partial X}{\partial x^k}(\pi_{x^0}(t)) + \frac{\partial \eta_k}{\partial t^\gamma} \frac{\partial X}{\partial x_\gamma^k}(\pi_{x^0}(t)) \right\} dv(<) &\leq -\rho b(x, x^0) \|\theta(x, x^0)\|^2. \end{aligned}$$

We underline that the notion of quasiinvexity is used in an appropriate form in a recent work by Nahak and Mohapatra [14] for a study of some multiobjective programming problems.

Having in mind the background, introduced above, now we can state and prove our results on duality [13], [22].

3. MOND-WEIR-ZALMAI TYPE DUALITY FOR (VFP)

Relative to (VFP), we consider the next multitime vector fractional variational dual problem

$$(ZMFD) \quad \begin{cases} \text{Maximize} & \left(\frac{F_1(y)}{K_1(y)}, \dots, \frac{F_p(y)}{K_p(y)} \right) \\ \text{subject to} & \tau^r \left[K_r(y) \frac{\partial f_r}{\partial x^k} - F_r(y) \frac{\partial k_r}{\partial x^k} \right] + \lambda^\alpha(t) \frac{\partial g_\alpha}{\partial x^k} + \mu^\beta(t) \frac{\partial h_\beta}{\partial x^k} \\ & - \frac{\partial}{\partial t^\gamma} \left(\tau^r \left[K_r(x) \frac{\partial f_r}{\partial x_\gamma^k} - F_r(x) \frac{\partial k_r}{\partial x_\gamma^k} \right] + \lambda^\alpha(t) \frac{\partial g_\alpha}{\partial x_\gamma^k} + \mu^\beta(t) \frac{\partial h_\beta}{\partial x_\gamma^k} \right) = 0, \\ & \lambda^\alpha(t) g_\alpha(\pi_y(t)) + \mu^\beta(t) h_\beta(\pi_y(t)) \geq 0, \quad i = \overline{1, p}, \quad \alpha = \overline{1, m}, \quad \beta = \overline{1, q}, \\ & \tau = (\tau^r) \geq 0, \quad e' \tau = 1, \quad (\mu^\beta(t)) \geq 0, \quad y \in \Phi, \quad y(t)|_{\partial\Omega} = u(t), \quad t \in \Omega. \end{cases}$$

Denote by $\varpi(x)$ the value of problem (VFP) at $x \in \mathcal{D}$ and be $\delta(y, \lambda, \eta, \nu)$ the value of the dual (ZMFD) at $(y, \lambda, \eta, \nu) \in \Delta$, where Δ is the domain of (ZMFD). In what follows we develop a duality theory between (VFP) and (ZMFD).

Theorem 3.1 (Weak duality). *Let x and (y, τ, λ, μ) be feasible points of problems (VFP) and (ZMFD), respectively. Assume that there are satisfied the next conditions:*

- a) for each $r = \overline{1, p}$ we have $F_r(x) > 0, K_r(x) > 0, \forall x \in X$;
 - b) for each $r = \overline{1, p}$, $F_r(x)$ is (ρ'_r, b) -quasiinvex at y and $-K_r(x)$ is (ρ''_r, b) -quasiinvex at y , all in respect to η and θ ;
 - c) $\int_{\Omega} \lambda^\alpha(t) g_\alpha(\pi_x(t)) dv$ is (ρ'''_r, b) -quasiinvex at y with respect to η and θ ;
 - d) $\int_{\Omega} \mu^\beta(t) h_\beta(\pi_x(t)) dv$ is monotonic (ρ^4, b) -quasiinvex at y with respect to η and θ ;
 - e) one of the functions of b) and c) is strictly (ρ, b) -quasiinvex at y with respect to η and θ ;
 - f) $\tau^r [\rho'_r K_r(y) + \rho''_r F_r(y)] + \rho'''_r + \rho^4 \geq 0$.
- Then $\varpi(x) \leq \delta(y, \tau, \lambda, \mu)$ is false.

Proof. From the first assumption of b), it follows

$$\begin{aligned} F_r(x) \leq F_r(y) &\Rightarrow \\ b(x, y) \int_{\Omega} \left\{ \eta_k \frac{\partial f_r}{\partial y^k} + \frac{\partial \eta_k}{\partial t^\gamma} \frac{\partial f_r}{\partial y_\gamma^k} \right\} dv &\leq -\rho'_r b(x, y) \|\theta(x, y)\|^2, \end{aligned} \quad (3.1)$$

while from the second assumption, we get

$$\begin{aligned} -K_r(x) \leq -K_r(y) &\Rightarrow \\ b(x, y) \int_{\Omega} \left\{ -\eta_k \frac{\partial k_r}{\partial y^k} - \frac{\partial \eta_k}{\partial t^\gamma} \frac{\partial k_r}{\partial y_\gamma^k} \right\} dv &\leq -\rho''_r b(x, y) \int_a^b \|\theta(x, y)\|^2. \end{aligned} \quad (3.2)$$

Multiplying relation (3.1) by $K_r(y) > 0$ and (3.2) by $F_r(y) > 0$, and summing the obtained implications side by side we get

$$\begin{aligned} F_r(x)K_r(y) - K_r(x)F_r(y) &\leq 0 \Rightarrow \\ b(x, y) \int_{\Omega} \left\{ \eta_k \left[K_r(y) \frac{\partial f_r}{\partial y^k} - F_r(y) \frac{\partial k_r}{\partial y^k} \right] + \frac{\partial \eta_k}{\partial t^\gamma} \left[K_r(y) \frac{\partial f_r}{\partial y_\gamma^k} - F_r(y) \frac{\partial k_r}{\partial y_\gamma^k} \right] \right\} dv & \quad (3.3) \\ &\leq -[\rho'_r K_r(y) + \rho''_r F_r(y)] b(x, y) \|\theta(x, y)\|^2. \end{aligned}$$

Multiplying (3.3) by τ^r , $r = \overline{1, p}$ ($\tau \geq 0$) and by summing over $r = \overline{1, p}$, we obtain

$$\begin{aligned} \tau^r [F_r(x)K_r(y) - K_r(x)F_r(y)] &\leq 0 \Rightarrow \\ b(x, y) \int_a^b \left\{ \eta_k \tau^r \left[K_r(y) \frac{\partial f_r}{\partial y^k} - F_r(y) \frac{\partial k_r}{\partial y^k} \right] + \frac{\partial \eta_k}{\partial t^\gamma} \tau^r \left[K_r(y) \frac{\partial f_r}{\partial y_\gamma^k} - F_r(y) \frac{\partial k_r}{\partial y_\gamma^k} \right] \right\} dv & \quad (3.4) \\ &\leq -b(x, y) \|\theta(x, y)\|^2 \tau^r [\rho'_r K_r(y) + \rho''_r F_r(y)]. \end{aligned}$$

From c) and d) respectively we have

$$\begin{aligned} \int_{\Omega} \lambda^\alpha(t) g_\alpha(\pi_x(t)) dv &\leq \int_{\Omega} \lambda^\alpha(t) g_\alpha(\pi_y(t)) dv \Rightarrow \\ b(x, y) \int_{\Omega} \left\{ \eta_k \left[\lambda^\alpha(t)' \frac{\partial g_\alpha}{\partial y^k} \right] + \frac{\partial \eta_k}{\partial t^\gamma} \left[\lambda^\alpha(t) \frac{\partial g_\alpha}{\partial y_\gamma^k} \right] \right\} dv &\leq -\rho'' b(x, y) \|\theta\|^2; \end{aligned} \quad (3.5)$$

and

$$\begin{aligned} \int_{\Omega} \mu^\beta(t) h_\beta(\pi_x(t)) dv &\leq \int_{\Omega} \mu^\beta(t) h_\beta(\pi_y(t)) dv \Rightarrow \\ b(x, y) \int_{\Omega} \left\{ \eta_k \left[\mu^\beta(t) \frac{\partial h_\beta}{\partial y^k} \right] + \frac{\partial \eta_k}{\partial t^\gamma} \left[\mu^\beta(t) \frac{\partial h_\beta}{\partial y_\gamma^k} \right] \right\} dv &\leq -\rho''' b(x, y) \|\theta\|^2. \end{aligned} \quad (3.6)$$

Summing now side by side the double implications (3.4), (3.5) and (3.6), and taking into account e) it follows

$$\begin{aligned} \tau^r [F_r(x)K_r(y) - K_r(x)F_r(y)] + \int_{\Omega} [\lambda^\alpha(t) g_\alpha(\pi_x(t)) + \mu^\beta(t) h_\beta(\pi_x(t))] dv & \\ - \int_{\Omega} [\lambda^\alpha(t) g_\alpha(\pi_y(t)) + \mu^\beta(t) h_\beta(\pi_y(t))] dv &\leq 0 \Rightarrow \\ b(x, y) \int_{\Omega} \left\{ \eta_k \left\{ \tau^r \left[K_r(y) \frac{\partial f_r}{\partial y^k} - F_r(y) \frac{\partial k_r}{\partial y^k} \right] + \lambda^\alpha(t)' \frac{\partial g_\alpha}{\partial y^k} + \mu^\beta(t) \frac{\partial h_\beta}{\partial y^k} \right\} \right. & \quad (3.7) \\ \left. + \frac{\partial \eta_k}{\partial t^\gamma} \tau^r \left[K_r(y) \frac{\partial f_r}{\partial y_\gamma^k} - F_r(y) \frac{\partial k_r}{\partial y_\gamma^k} \right] + \lambda^\alpha(t) \frac{\partial g_\alpha}{\partial y_\gamma^k} + \mu^\beta(t) \frac{\partial h_\beta}{\partial y_\gamma^k} \right\} dv & \\ < -b(x, y) \|\theta(x, y)\|^2 \{ \tau^r [\rho'_r K_r(y) + \rho''_r F_r(y)] + \rho''' + \rho^4 \}. & \end{aligned}$$

From the second implication of (3.7), we obtain $b(x, y) > 0$.

Then the second implication of (3.7) shortly, becomes

$$\int_{\Omega} \left[\eta_s \frac{\partial V}{\partial y^s} + \frac{\partial \eta_s}{\partial t^\gamma} \frac{\partial V}{\partial y_\gamma^s} \right] dt < -\|\theta(x, y)\|^2 \{ \tau^r [\rho_r' K_r(y) + \rho_r'' F_r(y)] + \rho''' + \rho^4 \}, \quad (3.8)$$

with

$$V = \tau^r [K_r(y) f_r(\pi_y(t)) - F_r(y) k_r(\pi_y(t))] + \lambda^\alpha(t) g_\alpha(\pi_y(t)) + \mu^\beta(t) h_\beta(\pi_y(t)).$$

For each index j , we have

$$\frac{\partial \eta_k}{\partial t^\gamma} \frac{\partial V}{\partial y_\gamma^k} = \frac{\partial}{\partial t^\gamma} \left(\eta_k \frac{\partial V}{\partial y_\gamma^k} \right) - \eta_k \frac{\partial}{\partial t^\gamma} \left(\frac{\partial V}{\partial y_\gamma^k} \right),$$

and, by integration, we obtain

$$\int_{\Omega} \frac{\partial \eta_k}{\partial t_j} \frac{\partial V}{\partial y_\gamma^k} dv = \int_{\Omega} \frac{\partial}{\partial t^\gamma} \left(\eta_k \frac{\partial V}{\partial y_\gamma^k} \right) dv - \int_{\Omega} \eta_k \frac{\partial}{\partial t^\gamma} \left(\frac{\partial V}{\partial y_\gamma^k} \right) dv.$$

Using the gradient formula, we get

$$\int_{\Omega} \frac{\partial}{\partial t^\gamma} \left(\eta_k \frac{\partial V}{\partial y_\gamma^k} \right) dv = \int_{\partial\Omega} \left(\eta_k \frac{\partial V}{\partial y_\gamma^k} \right) \vec{n}(t) d\sigma = 0,$$

where $\vec{n}(t)$ is unit vector to surface $\partial\Omega$ at the current point, and $\eta_s(t)|_{\partial\Omega} = 0$.

Then relation (3.8) becomes

$$\int_{\Omega} \eta_k \left[\frac{\partial V}{\partial y^k} - \frac{\partial}{\partial t^\gamma} \left(\frac{\partial V}{\partial y_\gamma^k} \right) \right] dv < -\|\theta(x, y)\|^2 \{ \tau^r [\rho_r' K_r(y) + \rho_r'' F_r(y)] + \rho''' + \rho^4 \}. \quad (3.9)$$

Taking into account the first constraint of problem (ZMFD), we have

$$\frac{\partial V}{\partial y^k} - \frac{\partial}{\partial t^\gamma} \left(\frac{\partial V}{\partial y_\gamma^k} \right) = 0$$

and relation (3.9) becomes

$$0 < -\|\theta(x, y)\|^2 \{ \tau^r [\rho_r' K_r(y) + \rho_r'' F_r(y)] + \rho''' + \rho^4 \}.$$

According to hypothesis f), this inequality becomes $0 < 0$ which is false. Then, from (3.7) is true the next relation

$$\begin{aligned} & \tau^r [F_r(x) K_r(y) - K_r(x) F_r(y)] + \int_{\Omega} [\lambda^\alpha(t) g_\alpha(\pi_x(t)) + \mu^\beta(t) h_\beta(\pi_x(t))] dv \\ & - \int_{\Omega} [\lambda^\alpha(t) g_\alpha(\pi_y(t)) + \mu^\beta(t) h_\beta(\pi_y(t))] dv > 0. \end{aligned} \quad (3.10)$$

Taking into account relations (3.5) and (3.6), from (3.10) it follows

$$\tau^r [F_r(x) K_r(y) - K_r(x) F_r(y)] > 0,$$

that contradicts relation (3.3). Therefore, $\varpi(x) \leq \delta(y, \tau, \lambda, \mu)$ is false. □

Theorem 3.2 (Direct duality). *Let x^0 be a normal efficient solution of primal (VFP) and suppose satisfied the hypotheses of Theorem 3.1. Then there are scalar $\tau^0 \in \mathbb{R}^p$ and the piecewise smooth functions $\lambda^0: \Omega \rightarrow \mathbb{R}^m$ and $\mu^0: \Omega \rightarrow \mathbb{R}^q$ such that $(x^0, \tau^0, \lambda^0, \mu^0)$ is an efficient solution to dual (ZMFD) and $\varpi(x^0) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.*

Proof. Since x^0 is a normal efficient solution to problem (VFP), according to Theorem 2.2 there are vector $\tau^0 = (\tau^r)^0 \in \mathbb{R}^p$ and the piecewise smooth functions $\lambda^0 = (\lambda^\alpha)^0: \Omega \rightarrow \mathbb{R}^{nm}$ and $\mu^0 = (\mu^\beta)^0: \Omega \rightarrow \mathbb{R}^q$ that satisfy relations (MFJ). We get $\mu^\beta(t)h_\beta(\pi_{x^0}(t)) = 0$, $\beta = \overline{1, q}$. Also $(\lambda^\alpha)^0(t)'g_\alpha(\pi_{x^0}(t)) = 0$ is true for all i and α . It follows that $(x^0, \tau^0, \lambda^0, \mu^0) \in \Delta$ and $\varpi(x^0) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$. \square

Theorem 3.3 (Converse duality). *Let $(x^0, \tau^0, \lambda^0, \mu^0)$ be an efficient solution of dual (MVFD) and suppose satisfied the following conditions:*

- i) \bar{x} is a normal efficient solution of primal (VFP);
 - ii) the hypotheses of Theorem 3.1 are satisfied for $(y, \tau, \lambda, \mu) = (x^0, \tau^0, \lambda^0, \mu^0)$.
- Then $\bar{x} = x^0$ and moreover, $\varpi(x^0) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.*

Proof. On the contrary, suppose that $\bar{x} \neq x^0$ and we shall obtain a contradiction. Because \bar{x} is a normal efficient solution to (VFP) then, according to Theorem 2.2 there are vector $\bar{\tau} \in \mathbb{R}^p$ and vector functions $\bar{\lambda} = (\bar{\lambda}^\alpha): \Omega \rightarrow \mathbb{R}^{nm}$ and $\bar{\mu} = (\bar{\mu}^\beta): \Omega \rightarrow \mathbb{R}^q$ that satisfy conditions (MFJ). We obtain

$$\bar{\lambda}^\alpha(t)g_\alpha(\pi_{\bar{x}}(t)) + \bar{\mu}^\beta(t)h_\beta(\pi_{\bar{x}}(t)) = 0$$

and so, $(\bar{x}, \bar{\tau}, \bar{\lambda}, \bar{\mu}) \in \Delta$. Moreover, $\varpi(\bar{x}) = \delta(\bar{x}, \bar{\tau}, \bar{\lambda}, \bar{\mu})$. According to Theorem 3.1 relation $\varpi(\bar{x}) \leq \delta(x^0, \tau^0, \lambda^0, \mu^0)$ is false. It results that the relation

$$\delta(\bar{x}, \bar{\tau}, \bar{\lambda}, \bar{\mu}) \leq \delta(x^0, \tau^0, \lambda^0, \mu^0),$$

is false. Therefore, the maximal efficiency of $(x^0, \tau^0, \lambda^0, \mu^0)$ is contradicted. Then, it gets that the supposition $\bar{x} \neq x^0$, above made, is false. It follows $\bar{x} = x^0$ and also $\bar{\tau} = \tau^0$, $\bar{\lambda} = \lambda^0$, $\bar{\mu} = \mu^0$. Finally, we obtain $\varpi(\bar{x}) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$. \square

Corollary 3.1 (Weak duality). *Let $x \in \mathcal{D}$ and $(y, \tau, \lambda, \mu) \in \Delta$ be feasible points of problems (VFP) and (ZMFD). Assume that are satisfied the conditions:*

- a) for each $r = \overline{1, p}$, integral $\int_{\Omega} [K_r(y)f_r(\pi_x(t)) - F_r(y)k_r(\pi_x(t))]dv$ is (ρ'_r, b) -quasiinvex at y with respect to η and θ ;
 - b) $\int_{\Omega} [\lambda^\alpha(t)g_\alpha(\pi_x(t)) + \mu^\beta(t)h_\beta(\pi_x(t))]dv$ is (ρ, b) -quasiinvex at y with respect to η and θ ;
 - c) one of the functions of a)-b) is strictly (ρ, b) -quasiinvex at y with respect to η and θ ;
 - d) $\tau^r \rho'_r + \rho \geq 0$.
- Then $\varpi(x) \leq \delta(y, \tau, \lambda, \mu)$ is false.*

Corollary 3.2 (Direct duality). *Let x^0 be a normal efficient solution of the primal (VFP) and suppose satisfied the hypotheses of Theorem 3.1. Then there are vector*

$\tau^0 \in \mathbb{R}^p$ and the piecewise smooth functions $\lambda^0 = (\lambda^\alpha)^0: \Omega \rightarrow \mathbb{R}^{nm}$ and $\mu^0 = (\mu^\alpha)^0: \Omega \rightarrow \mathbb{R}^q$ such that $(x^0, \tau^0, \lambda^0, \mu^0)$ is an efficient solution of the dual (ZMFD) and $\varpi(x^0) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.

Corollary 3.3 (Converse duality). *Let $(x^0, \tau^0, \lambda^0, \mu^0) \in \Delta$ be a efficient solution to the dual and (ZMFD) and assume satisfied the next conditions:*

- i) \bar{x} is a normal efficient solution of the primal (VFP);
- ii) the hypotheses of Theorem 3.1 are satisfied with $(y, \tau, \lambda, \mu) = (x^0, \tau^0, \lambda^0, \mu^0)$.
Then $\bar{x} = x^0$ and $\varpi(x^0) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.

The theory in this section allows to consider some special cases of interest. These ones make the object of the following two sections.

4. CASE STUDY 1: MOND-WEIR DUALITY

In this section, we consider $K_r(x) = 1$, $r = \overline{1, p}$. Problem (VFP) becomes (VVP), and (ZMFD) becomes the following multitime vector variational problem :

$$(MWD) \begin{cases} \text{Maximize } \int_{\Omega} f(\pi_y(t))dv = (F_1(y), \dots, F_p(y)) \\ \text{subject to } \tau^r \frac{\partial f_r}{\partial y^k} + \lambda^\alpha(t) \frac{\partial g_\alpha}{\partial y^k} + \mu^\beta(t) \frac{\partial h_\beta}{\partial y^k} \\ \qquad \qquad \qquad - \frac{\partial}{\partial t^\gamma} \left(\tau^i \frac{\partial f_i}{\partial y^k_\gamma} + \lambda^\alpha(t) \frac{\partial g_\alpha}{\partial y^k_\gamma} + \mu^\beta(t) \frac{\partial h_\beta}{\partial y^k_\gamma} \right) = 0 \\ \lambda^\alpha(t)' g_\alpha(\pi_y(t)) + \mu^\beta(t) h_\beta(\pi_y(t)) \geq 0 \text{ [no summation]} \\ \tau \geq 0, e'\tau = 1, y(t)|_{\partial\Omega} = u(t), t \in \Omega. \end{cases}$$

The domain of (VVP) is \mathcal{D} and denote by Δ' the domain of (MWD). Denote by $\varpi(x)$ the value of problem (VVP) at $x \in \mathcal{D}$ and be $\delta(y, \tau, \lambda, \mu)$ the value of the dual (MWD) at $(y, \tau, \lambda, \mu) \in \Delta'$.

There exist the following duality theorems of Mond-Weir type between problems (VVP) and (MWD).

Theorem 4.1 (Weak duality). *Let $x \in \mathcal{D}$ and $(y, \tau, \lambda, \mu) \in \Delta$ be feasible points of the problems (VVP) and (MWD) respectively. Assume the conditions hold:*

- a) for each $r = \overline{1, p}$, the integral $\int_{\Omega} f_r(\pi_x(t))dv$ is (ρ'_r, b) -quasiinvex at $x = y$ with respect to η and θ ;
- b) $\int_{\Omega} \lambda^\alpha(t) g_\alpha(\pi_x(t))dv$ [no summation] is (ρ''_α, b) -quasiinvex at y with respect to η and θ ;
- c) $\int_{\Omega} \mu^\beta(t) h_\beta(\pi_x(t))dv$ [no summation] is monotonic (ρ'''_β, b) -quasiinvex at y with respect to η and θ ;

d) one of the functions of a)-c) is strictly (ρ, b) -quasiinvex at y with respect to η and θ ;

$$e) \tau^r \rho'_r + \sum_{\alpha=1}^m \rho''_{\alpha} + \sum_{\beta=1}^q \rho'''_{\beta} \geq 0.$$

Then $\varpi(x) \leq \delta(y, \tau, \lambda, \mu)$ is false.

Theorem 4.2 (Direct duality). *Let x^0 be a normal efficient solution of the primal (VVP) and suppose satisfied the hypotheses of Theorem 4.1. Then there are $\tau^0 \in \mathbb{R}^p$ and the piecewise smooth functions $\lambda^0: \Omega \rightarrow \mathbb{R}^m$ and $\mu^0: \Omega \rightarrow \mathbb{R}^q$ such that $(x^0, \tau^0, \lambda^0, \mu^0)$ is an efficient solution of the dual problem (MWD). Moreover, $\varpi(x^0) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.*

Theorem 4.3 (Converse duality). *Let $(x^0, \tau^0, \lambda^0, \mu^0)$ be an efficient solution of the dual (MWD) and suppose satisfied the following conditions:*

- i) \bar{x} is a normal efficient solution of the primal (VVP);
 - ii) the hypotheses of Theorem 4.1 are satisfied for $(y, \tau, \lambda, \mu) = (x^0, \tau^0, \lambda^0, \mu^0)$.
- Then $\bar{x} = x^0$ and $\varpi(x^0) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.

As we expected, Theorems 4.1-4.3 are particular cases of Theorems 3.1-3.3.

Corollary 4.1 (Weak duality). *Let x and (y, τ, λ, μ) be feasible points of problems (VVP) and (MWD). Assume that are satisfied the conditions:*

- a) for each $r = \overline{1, p}$, the integral $\int_{\Omega} f_r(\pi_x(t)) dv$ is (ρ'_r, b) -quasiinvex at y with respect to η and θ ;
 - b) $\int_{\Omega} [\lambda^{\alpha}(t)g_{\alpha}(\pi_x(t)) + \mu^{\beta}(t)h_{\beta}(\pi_x(t))] dv$ is (ρ, b) -quasiinvex at y with respect to η and θ ;
 - c) one of the functions of a)-b) is strictly quasiinvex at y with respect to η and θ ;
 - d) $\tau^r \rho'_r + \rho \geq 0$.
- Then $\varpi(x) \leq \delta(y, \lambda, \mu, v)$ is false.

Corollary 4.2 (Direct duality). *Let x^0 be a normal efficient solution of primal (VVP) and suppose satisfied the hypotheses stated in Corollary 4.1. There are vector $\tau^0 \in \mathbb{R}^p$ and the piecewise smooth functions $\lambda^0 = (\lambda^{\alpha})^0: \Omega \rightarrow \mathbb{R}^{nm}$ and $\mu^0 = (\mu^{\beta})^0: \Omega \rightarrow \mathbb{R}^q$ such that $(x^0, \tau^0, \lambda^0, \mu^0)$ is an efficient solution of the dual (MWD). Moreover, $\varpi(x^0) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.*

Corollary 4.3 (Converse duality). *Let $(x^0, \tau^0, \lambda^0, \mu^0)$ be an efficient solution to the dual (MWD) and suppose satisfied the following conditions:*

- i) \bar{x} is a normal efficient solution of primal (VP);
 - ii) the hypotheses of Corollary 4.1 are satisfied for $(y, \tau, \lambda, \mu) = (x^0, \tau^0, \lambda^0, \mu^0)$.
- Then $\bar{x} = x^0$ and $\varpi(\bar{x}) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.

Corollaries 4.1-4.3 are particular cases of Corollaries 3.1-3.3.

5. CASE STUDY 2: WOLFE DUALITY

Dual problem of Wolfe type, associated to (VVP) is the next multitime vector variational problem of maximum

$$(WD) \quad \begin{cases} \text{Maximize } \int_{\Omega} \{f(\pi_y(t)) + [\lambda^\alpha(t)g_\alpha(\pi_y(t)) + \mu^\beta(t)h_\beta(\pi_y(t))]\} dv \\ \text{subject to } \tau^r \frac{\partial f_r}{\partial y^k} + \lambda^\alpha(t) \frac{\partial g_\alpha}{\partial y^k} + \mu^\beta(t) \frac{\partial h_\beta}{\partial y^k} - \\ \quad - \frac{\partial}{\partial t^\gamma} \left(\tau^r \frac{\partial f_r}{\partial y_\gamma^k} + \lambda^\alpha(t) \frac{\partial g_\alpha}{\partial y_\gamma^k} + \mu^\beta(t) \frac{\partial h_\beta}{\partial y_\gamma^k} \right) = 0, \\ \mu^\beta(t) \geq 0, \forall t \in \Omega, k = \overline{1, n}, j = \overline{1, m}, e'\tau = 1. \end{cases}$$

Corollary 5.1 (Weak duality). *Let x and (y, τ, λ, μ) be feasible points of problems (VVP) and (WD). Assume that are satisfied the conditions:*

- a) *for each $r = \overline{1, p}$, the integral $\int_{\Omega} f_r(\pi_x(t)) dv$ is (ρ'_r, b) -quasiinvex at y with respect to η and θ ;*
- b) *$\int_{\Omega} [\lambda^\alpha(t)g_\alpha(\pi_x(t)) + \mu^\beta(t)h_\beta(\pi_x(t))] dv$ is (ρ, b) -quasiinvex at y with respect to η and θ ;*
- c) *one of the functions of a)-b) is strictly quasiinvex at y with respect to η and θ ;*
- d) *$\tau^r \rho'_r + \rho \geq 0$.*

Then $\varpi(x) \leq \delta(y, \tau, \lambda, \mu)$ is false.

Corollary 5.2 (Direct duality). *Let x^0 be a normal efficient solution of primal (VVP) and suppose satisfied the hypotheses stated in Corollary 5.1. Then there are vector $\tau^0 \in \mathbb{R}^p$ and the piecewise smooth functions $\lambda^0 = (\lambda^\alpha)^0: \Omega \rightarrow \mathbb{R}^{nm}$ and $\mu^0 = (\mu^\beta)^0: \Omega \rightarrow \mathbb{R}^q$ such that $(x^0, \tau^0, \lambda^0, \mu^0)$ is an efficient solution of dual (WD) and moreover, $\varpi(x^0) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.*

Corollary 5.3 (Converse duality). *Let $(x^0, \tau^0, \lambda^0, \mu^0)$ be an efficient solution to the dual (DWP) and suppose satisfied the following conditions:*

- i) *\bar{x} is a normal efficient solution of the primal (VP);*
- ii) *the hypotheses of Corollary 4.1 are satisfied for $(y, \tau, \lambda, \mu) = (x^0, \tau^0, \lambda^0, \mu^0)$. Then $\bar{x} = x^0$ and moreover, $\varpi(\bar{x}) = \delta(x^0, \tau^0, \lambda^0, \mu^0)$.*

As we easily can see, Corollaries 5.1-5.3, stated above, are particular cases of Corollaries 4.1-4.3.

6. CONCLUSION

In this paper, we considered a study of a multitime vector (or multiobjective) variational problem (VVP) and a multitime vector fractional variational problem (VFP). For the two vector variational problems, Mititelu and Postolache established necessary efficiency conditions in their joint research work [11]. It was the aim of

the present paper to utilize these conditions, in order to establish new duality conditions of Mond-Weir-Zalmai type for the fractional problem (VFP) through weak, direct and converse duality theorems. As particular cases, we obtained some duality conditions for (VVP). The notion of (ρ, b) -quasiinvexity was the main ingredient in the development of our theory.

The present study, was motivated by the wide class of applications of fractional programming arising from applied areas including engineering design, game theory, stock cutting, portfolio selection, decision problems in management. For a review on theory and applications of multiobjective programming, we address the reader to the monograph by Jahn, [5].

REFERENCES

- [1] A. Chinchuluun and P. M. Pardalos: *A survey of recent developments in multiobjective optimization*, Ann. Oper. Res., **154**(2007), 29-50.
- [2] M. A. Geoffrion: *Proper efficiency and the theory of vector minimization*, J. Math. Anal. Appl., **149**(1968), No. 6, 618-630.
- [3] M. A. Hanson: *Duality for variational problems*, J. Math. Anal. Appl., **18**(1967), 355-364.
- [4] R. Jagannathan: *Duality for nonlinear fractional programming*, Z. Oper. Res., **17**(1973), 1-3.
- [5] J. Jahn: *Vector Optimization. Theory, Applications and Extensions*, Springer, Berlin, 2004.
- [6] V. Jeyakumar and B. Mond: *On generalized convex mathematical programming*, J. Austral. Math. Soc., Series B, **34**(1992), 4353.
- [7] Z. Khan and M. A. Hanson: *On ratio invexity in mathematical programming*, J. Math. Anal. Appl., **205**(1997), 330-336.
- [8] Z. A. Liang, H. X. Huang and P. M. Pardalos: *Efficiency conditions and duality for a class of multiobjective fractional programming problems*, J. Glob. Optim., **27**(2003), 447-471.
- [9] Șt. Mititelu: *Efficiency conditions for multiobjective fractional variational problems*, Appl. Sci., **10**(2008), 162-175.
- [10] Șt. Mititelu: *Optimality and duality for invex multi-time control problems with mixed constraints*, J. Adv. Math. Stud., **2**(2009), No. 1, 25-34.
- [11] Șt. Mititelu and M. Postolache: *Efficiency and duality for multitime vector fractional variational problems on manifolds*, Balkan J. Geom. Appl., **16**(2011), No. 2, 90-101.
- [12] Șt. Mititelu and I. M. Stancu-Minasian: *Efficiency and duality for multiobjective fractional variational problems with (ρ, b) -quasiinvexity*, Yugoslav J. Oper. Res., **19**(2009), No. 1, 85-99.
- [13] B. Mond and T. Weir: *Generalized concavity and duality*, in Generalized Concavity in Optimization and Economics (S. Schaible and W. T. Ziemba (Eds.)), Proc. NATO Adv. Study Inst., Vancouver/Can. 1980, 263-279 (1981).
- [14] C. Nahak and R. N. Mohapatra: *Nonsmooth ρ - (η, θ) -invexity in multipbjective programming problems*, Optim. Lett., DOI 10.1007/s11590-010-0239-1.
- [15] Ariana Pitea, C. Udriște and Șt. Mititelu: *PDE constrained optimization problems with curvilinear functional quotiens as objective vectors*, Balkan J. Geom. Appl., **14**(2009), No. 2, 75-88.
- [16] Ariana Pitea, C. Udriște and Șt. Mititelu: *New type of duality in PDI and PDE constrained optimization problems*, J. Adv. Math. Stud., **2**(2009), No. 1, 75-88.

- [17] V. Preda: *On efficiency and duality for multiobjective programs*, J. Math. Anal. Appl., **166**(1992), No. 2, 365-377.
- [18] V. Preda: *On Mond-Weir duality for variational problems*, Rev. Roumanie Math. Pures Appl., **28**(1993), No. 2, 155-164.
- [19] L. V. Reddy and R. N. Mukherjee: *Some results on mathematical programming with generalized ratio invexity*, J. Math. Anal. Appl., **240**(1999), 299-310.
- [20] C. Singh and M. A. Hanson: *Multiobjective fractional programming duality theory*, Naval Research Logistics, **38**(1991), 925-933.
- [21] C. Udriște: *Simplified multitime maximum principle*, Balkan J. Geom. Appl., **14**(2009), No. 1, 102-119.
- [22] C. Udriște and I. Tevy: *Multitime Euler-Lagrange-Hamilton theory*, WSEAS Trans. Math., **6**(2007), No. 6, 701-709.
- [23] F. A. Valentine: *The problem of Lagrange with differentiable inequality as added side conditions*, 407-448, in „Contributions to the Calculus of Variations”, 1933-37, Univ. of Chicago Press, 1937.
- [24] G. J. Zalmai: *Generalized $(\mathcal{F}, b, \varphi, \rho, \theta)$ -univex n -set functions and semiparametric duality models in multiobjective fractional subset programming*, Int. J. Math. Math. Sci., 2005, No. 7, 1109-1133.

*Technical University of Civil Engineering
Department of Mathematics and Informatics
Bd. Lacul Tei, No. 124, 020396 Bucharest, Romania
E-mail address: st.mititelu@yahoo.com*

*University of Bucharest
Faculty of Mathematics and Informatics
Academiei, No. 14, 70109 Bucharest, Romania
E-mail address: preda@fmi.mathem.ro*

*University “Politehnica” of Bucharest
Faculty of Applied Sciences
Splaiul Independenței, No. 313, 060042 Bucharest, Romania
E-mail address: mihai@mathem.pub.ro*